



CPAL

Central Pennsylvania Alliance Laboratory

Technical Bulletin

No. 58

May 16, 2006

Factor II (Prothrombin) 20210A Genetic Analysis

Starting Date:

Currently Available

Mnemonic: PROTH MUT

Contact:

Dr. Jeffrey Wisotzkey, 717-851-1422
Technical Director, CPAL

Summary:

The Factor II (Prothrombin) G20210A assay allows the detection and genotyping of a single point mutation (G to A at position 20210) of the human Factor II gene from DNA isolated from human whole peripheral blood. The test is performed on the LightCycler instrument utilizing polymerase chain reaction (PCR) for the amplification of Factor II DNA recovered from clinical specimens and fluorogenic target-specific hybridization for the detection and genotyping of the amplified Factor II DNA.

Detection and genotyping of the Factor II (Prothrombin) G20210A mutation aids in the evaluation of patients with suspected thrombophilia. Inherited thrombophilia predispose an individual to thrombotic events such as venous thrombosis, the third most common cardiovascular disease. Activated protein C (APC) resistance is regarded as the most prevalent coagulation abnormality associated with venous thrombosis. Patients tested positive for APC resistance or the Factor V Leiden mutation should be considered for molecular genetic testing for the most common other thrombophilias with overlapping phenotype, for which testing is available at present [*i.e.*, the Factor II (Prothrombin) G20210A variant]. It is present in 1-2% of the general population and its involvement in venous thromboembolism is well established.

Reference Ranges:

The “normal” result for this assay is reported as WILDTYPE. This indicates that the mutation, better termed “polymorphism”, is not present on either of the two copies of the factor II gene (allele).

Issued on: May 16, 2006

For questions about this and other information, call Central Pennsylvania Alliance Laboratory at 1-888-480-1422.

Important Additional Notes:

- This assay is a germline genetic assay and should only be ordered once on any particular patient. The Prothrombin 20210A genotype is constitutional and does not change.
- The Prothrombin 20210A assay is typically ordered in conjunction with the Factor V Leiden mutational assay and other markers of hypercoagulability, in patients with thrombosis.

References:

1. Poort, SR, Rosendaal, FR, Reitsma, PH and Bertina, RM. *A common genetic variation in the 3'-untranslated region of the prothrombin gene is associated with elevated plasma prothrombin levels and an increase in venous thrombosis.* Blood. 1996 Nov 15;88(10):3698-703.
2. Wisotzkey, J.D., P. Bayliss, E. Rutherford and T. Bell. *Placental Genotyping of the Factor V Leiden, Prothrombin 20210A and the Methylenetetrahydrofolate Reductase (MTHFR) C677T Alleles in IUGR Pregnancies (Letter).* **Thrombosis and Haemostasis**, 81:844-845, 1999
3. Wisotzkey, J.D., Bell, T. and Monk, J.S. *Simultaneous Polymerase Chain Reaction Restriction Fragment Length Polymorphism (PCR-RFLP) Identification of the Factor V Leiden Allele and the Prothrombin 20210A Mutation.* **Diagnostic Molecular Pathology**, 7:180-183, 1998.